

Decay coupling constants sum rules for dibaryon octet into two baryon octets with λ_8 first order SU(3) symmetry breaking

E. N. Polanco-Euán,^{*} V. Gupta,[†] and G. Sánchez-Colón[‡]

Departamento de Física Aplicada.

Centro de Investigación y de Estudios Avanzados del IPN.

Unidad Mérida.

A.P. 73, Cordemex.

Mérida, Yucatán, 97310. MEXICO.

B. A. Bambah[§]

School of Physics.

University of Hyderabad.

Hyderabad, A.P. 500 046, India.

(Dated: July 14, 2016)

Abstract

The SU(3) octet states with baryon number $B = 2$, hexaquark dibaryons, are considered. Decay coupling constants sum rules for dibaryon octet into two ordinary baryon octets with λ_8 first order SU(3) symmetry breaking are given. An SU(4) extension of the analysis is commented upon. Possibilities for the experimental observation of multibaryon and anti-multibaryon states are pointed out.

PACS numbers: 14.20.Pt, 13.30.-a

Keywords: dibaryon, sextaquarks, sum rule

^{*}elias.polanco@mda.cinvestav.mx

[†]virendra@mda.cinvestav.mx

[‡]gabriel.sanchez@cinvestav.mx

[§]bbsp@uohyd.ernet.in

I. INTRODUCTION.

The first hexaquark or sextaquark state with baryon number $B = 2$, a dibaryon, was proposed in 1964 [1]. Hexaquarks received more attention after the proposal of the H-dibaryon ($uuddss$) by Jaffe [2]. The presence of hexaquark states have also been conjectured in models with colour flux tube (string) confinement, [3–5]. The theoretical case for the dibaryon (H-dibaryon) has recently been made stronger by lattice-QCD calculations in the SU(3) flavor limit by the NPLQCD and HALQCD collaborations [6, 7]. The presence of multi-quark states including the hexaquark in hot-dense matter is predicted in certain statistical [8] and holographic models [9], and since strangeness enhancement is associated with the QGP, hexaquarks with strangeness may have an observable signature in the upcoming heavy ion experiments at LHC. More recently Bashkanov *et. al* [10] have suggested novel dibaryon states and their possible experimental signatures. These include exclusive measurements of the $np \rightarrow np\pi^0\pi^0$ cross section at the WASA detector in COSY, Julich, which have shown the presence of a $d^*(2348)$ resonance that can be interpreted as a dibaryon state [11]. Recently in Ref. [12], it was noted that the deuteron, a stable dibaryon, belongs to the $B = 2$, $\mathbf{10}^*$ representation in the SU(3) $\mathbf{8} \otimes \mathbf{8}$ decomposition of two $J^P = (1/2)^+$ baryon octets (denoted here as B_8) and sum rules for decays of $\mathbf{10}^*$ and $\mathbf{10}$ dibaryons decuplets (denoted $D_{\mathbf{10}^*,\mathbf{10}}$) into an antisymmetric final state of two octet baryons, $D_{\mathbf{10}^*,\mathbf{10}} \rightarrow B_8 + B_8$, were given. A recent high statistics search in inclusive Upsilon decays was done for H-dibaryons in the symmetric SU(3) state at the Belle detector of KEKB e^+e^- collider [13], which put stringent constraints on the H-dibaryon mass and clearly showed that if a H-dibaryon exists in the mass range $\leq 2m_\Lambda$, then it must be different from a deuteron. M. Oka has also proposed the possibility of several dibaryon resonances examining flavor SU(3) octet dibaryon systems employing the quark cluster model [14]. Thus, it is important to study the various decay channels of a dibaryon belonging to the dibaryon octet.

In this paper we present sum rules for decays of the dibaryon SU(3) octet (denoted D_8) into two ordinary $J^P = (1/2)^+$ baryon octets with λ_8 first order SU(3) symmetry breaking. This additional study of the symmetry breaking mechanism at SU(3) level sets the stage for how to treat the breaking of SU(4) to get reasonable estimates for couplings of charmed hexaquark states.

In Sec. II the baryonic content of the D_8 states and their expected masses are presented.

In Sec. III decay sum rules in broken SU(3) are determined. In Sec. III A 1 the extension to the SU(4) case is considered.

Finally we discuss some open problems for multiquark states.

II. BARYONIC CONTENT AND MASSES OF DIBARYON OCTET

The (Y, I) states (with hypercharge Y and isospin I) of the dibaryon octet, denoted by $D_8(Y, I)$ are given in Table I. The last column gives an estimate of the average mass of $D_8(Y, I)$ depending on its baryonic content. From these we obtain:

$$3D_8(0, 0) + D_8(0, 1) = 9286.2\text{MeV}, \quad (1)$$

and

$$2[D_8(1, 1/2) + D_8(-1, 1/2)] = 9132.0\text{MeV}. \quad (2)$$

It is remarkable how well these estimate satisfy the first order Gell Mann-Okubo mass formula for the octet [17, 18].

III. SUM RULES FOR $D_8 \rightarrow B_8 + B_8$ DECAYS WITH FIRST ORDER SYMMETRY BREAKING OF SU(3).

We consider now dibaryon octet decays into a final state of two baryon octets: $D_8 \rightarrow B_8 + B_8$. Following Ref. [15] we write B_8 and D_8 matrices as:

$$B_8 = \begin{pmatrix} \frac{1}{\sqrt{6}}\Lambda + \frac{1}{\sqrt{2}}\Sigma^0 & \Sigma^+ & p \\ \Sigma^- & \frac{1}{\sqrt{6}}\Lambda - \frac{1}{\sqrt{2}}\Sigma^0 & n \\ -\Xi^- & \Xi^0 & -\sqrt{\frac{2}{3}}\Lambda \end{pmatrix}, \quad (3)$$

$$D_8 = \begin{pmatrix} \frac{1}{\sqrt{6}}D_8(0, 0, 0) + \frac{1}{\sqrt{2}}D_8(0, 1, 0) & D_8(0, 1, +1) & D_8(1, 1/2, +1/2) \\ D_8(0, 1, -1) & \frac{1}{\sqrt{6}}D_8(0, 0, 0) - \frac{1}{\sqrt{2}}D_8(0, 1, 0) & D_8(1, 1/2, -1/2) \\ -D_8(-1, 1/2, -1/2) & D_8(-1, 1/2, +1/2) & -\sqrt{\frac{2}{3}}D_8(0, 0, 0) \end{pmatrix}, \quad (4)$$

where the (Y, I, I_3) states of D_8 are denoted by $D_8(Y, I, I_3)$ with I_3 the isospin third component.

In general, for exact SU(3) the Yukawa interaction for $D_8 \rightarrow B_8 + B_8$ ($8 \rightarrow 8 \otimes 8$) is characterized by two parameters, g_8^0 and $g_{8'}^0$, since there are two octets in the decomposition:

$$8 \otimes 8 = 1 \oplus 8 \oplus 8' \oplus 10 \oplus 10^* \oplus 27. \quad (5)$$

The **1**, **8**, and **27** are symmetric under the interchange of the original two octets and **8'**, **10**, and **10*** are antisymmetric.

With λ_8 first order breaking of SU(3) we expect eight additional parameters in addition to g_8^0 and $g_{8'}^0$, since the one dimensional representation **1** appears eight times in the decomposition $8 \otimes 8 \otimes 8 \otimes 8$, four corresponding to couplings with flavor antisymmetric final state representations and four corresponding to couplings with flavor symmetric final state representations.

In this work we will analyze both cases, antisymmetric and symmetric final state, separately.

A. Decays of octet dibaryon into $B_8 + B_8$ antisymmetric final state.

We consider decays of the dibaryon octet into a flavor antisymmetric final state of two baryon octets. As described above, in this case we have five parameters: $g_{8'}^0$ from exact SU(3) and g_k ($k = 1, 2, 3, 4$) from λ_8 first order SU(3) symmetry breaking. The $D_8 \rightarrow B_8 + B_8$ Yukawa couplings in terms of $g_{8'}^0$ and g_k ($k = 1, 2, 3, 4$) are (for sake of clarity, we put a prime on the baryon matrix in the second octet and $\overline{D}_8 = D_8^\dagger$):

$$g_{8'}^0 \text{Tr} \{ \overline{D}_8 [B_8 B_8' - B_8' B_8] \}, \quad (6)$$

$$g_1 \text{Tr} \{ \overline{D}_8 \lambda_8 [B_8 B_8' - B_8' B_8] \}, \quad (7)$$

$$g_2 \text{Tr} \{ \overline{D}_8 [B_8 \lambda_8 B_8' - B_8' \lambda_8 B_8] \}, \quad (8)$$

$$g_3 \text{Tr} \{ \overline{D}_8 [B_8 B_8' - B_8' B_8] \lambda_8 \}, \quad (9)$$

$$g_4 \left\{ \text{Tr} [\overline{D}_8 B_8] \text{Tr} [B'_8 \lambda_8] - \text{Tr} [\overline{D}_{8_A} B'_8] \text{Tr} [B_8 \lambda_8] \right\}. \quad (10)$$

There are eight independent decay coupling constants to be determined. We take these eight constants to be:

$$G[D_8(1, 1/2, +1/2) \rightarrow p \Sigma^{0'}], \quad G[D_8(1, 1/2, +1/2) \rightarrow p \Lambda'], \quad (11)$$

$$G[D_8(0, 1, +1) \rightarrow p \Xi^{0'}], \quad G[D_8(0, 1, +1) \rightarrow \Sigma^+ \Lambda'], \quad G[D_8(0, 1, +1) \rightarrow \Sigma^+ \Sigma^{0'}], \quad (12)$$

$$G[D_8(0, 0, 0) \rightarrow p \Xi^{-'}], \quad (13)$$

$$G[D_8(-1, 1/2, +1/2) \rightarrow \Sigma^+ \Xi^{-'}], \quad G[D_8(-1, 1/2, +1/2) \rightarrow \Lambda \Xi^{0'}]. \quad (14)$$

The remaining decay coupling constants are given by:

$$\begin{aligned} \sqrt{2} G[D_8(1, 1/2, +1/2) \rightarrow p \Sigma^{0'}] &= -G[D_8(1, 1/2, +1/2) \rightarrow n \Sigma^{+'}] = \\ &= G[D_8(1, 1/2, -1/2) \rightarrow p \Sigma^{-'}] = -\sqrt{2} G[D_8(1, 1/2, -1/2) \rightarrow n \Sigma^{0'}], \end{aligned} \quad (15)$$

$$G[D_8(1, 1/2, +1/2) \rightarrow p \Lambda'] = G[D_8(1, 1/2, -1/2) \rightarrow n \Lambda'], \quad (16)$$

$$G[D_8(0, 1, +1) \rightarrow \Sigma^+ \Lambda'] = G[D_8(0, 1, 0) \rightarrow \Sigma^0 \Lambda'] = G[D_8(0, 1, -1) \rightarrow \Sigma^- \Lambda'], \quad (17)$$

$$\begin{aligned} G[D_8(0, 1, +1) \rightarrow p \Xi^{0'}] &= \sqrt{2} G[D_8(0, 1, 0) \rightarrow p \Xi^{-'}] = \sqrt{2} G[D_8(0, 1, 0) \rightarrow n \Xi^{0'}] = \\ &= G[D_8(0, 1, -1) \rightarrow n \Xi^{-'}], \end{aligned} \quad (18)$$

$$G[D_8(0, 1, +1) \rightarrow \Sigma^+ \Sigma^{0'}] = G[D_8(0, 1, 0) \rightarrow \Sigma^+ \Sigma^{-'}] = G[D_8(0, 1, -1) \rightarrow \Sigma^0 \Sigma^{-'}], \quad (19)$$

$$G[D_{\mathbf{8}}(0, 0, 0) \rightarrow n \Xi^{0'}] = -G[D_{\mathbf{8}}(0, 0, 0) \rightarrow p \Xi^{-'}], \quad (20)$$

$$\begin{aligned} G[D_{\mathbf{8}}(-1, 1/2, +1/2) \rightarrow \Sigma^+ \Xi^{-'}] &= -\sqrt{2} G[D_{\mathbf{8}}(-1, 1/2, +1/2) \rightarrow \Sigma^0 \Xi^{0'}] = \\ &= \sqrt{2} G[D_{\mathbf{8}}(-1, 1/2, -1/2) \rightarrow \Sigma^0 \Xi^{-'}] = -G[D_{\mathbf{8}}(-1, 1/2, -1/2) \rightarrow \Sigma^- \Xi^{0'}], \end{aligned} \quad (21)$$

$$G[D_{\mathbf{8}}(-1, 1/2, +1/2) \rightarrow \Lambda \Xi^{0'}] = G[D_{\mathbf{8}}(-1, 1/2, -1/2) \rightarrow \Lambda \Xi^{-'}]. \quad (22)$$

The eight independent decay coupling constants (11)-(14) are described by five parameters $g_{\mathbf{8}'}^0$ and g_k ($k = 1, 2, 3, 4$). Thus, three SU(3) broken sum rules between the Yukawa couplings may be deduced:

$$\begin{aligned} 4 G[D_{\mathbf{8}}(1, 1/2, +1/2) \rightarrow p \Sigma^{0'}] &= 2\sqrt{3} G[D_{\mathbf{8}}(0, 0, 0) \rightarrow p \Xi^{-'}] \\ &- \sqrt{2} G[D_{\mathbf{8}}(0, 1, +1) \rightarrow p \Xi^{0'}] + 2 G[D_{\mathbf{8}}(0, 1, +1) \rightarrow \Sigma^+ \Sigma^{0'}] \\ &- 2\sqrt{2} G[D_{\mathbf{8}}(-1, 1/2, +1/2) \rightarrow \Sigma^+ \Xi^{-'}], \end{aligned} \quad (23)$$

$$\begin{aligned} 2\sqrt{6} G[D_{\mathbf{8}}(1, 1/2, +1/2) \rightarrow p \Lambda'] &= 3\sqrt{6} G[D_{\mathbf{8}}(0, 0, 0) \rightarrow p \Xi^{-'}] \\ &+ 5 G[D_{\mathbf{8}}(0, 1, +1) \rightarrow p \Xi^{0'}] - \sqrt{2} G[D_{\mathbf{8}}(0, 1, +1) \rightarrow \Sigma^+ \Sigma^{0'}] \\ &+ 2\sqrt{6} G[D_{\mathbf{8}}(-1, 1/2, +1/2) \rightarrow \Xi^0 \Lambda'], \end{aligned} \quad (24)$$

$$\begin{aligned} \sqrt{6} G[D_{\mathbf{8}}(0, 1, +1) \rightarrow \Sigma^+ \Lambda'] &= \\ &= 2 G[D_{\mathbf{8}}(0, 1, +1) \rightarrow p \Xi^{0'}] - \sqrt{2} G[D_{\mathbf{8}}(0, 1, +1) \rightarrow \Sigma^+ \Sigma^{0'}] \\ &+ 3 G[D_{\mathbf{8}}(-1, 1/2, +1/2) \rightarrow \Sigma^+ \Xi^{-'}] + \sqrt{6} G[D_{\mathbf{8}}(-1, 1/2, +1/2) \rightarrow \Xi^0 \Lambda'], \end{aligned} \quad (25)$$

with identical relationships for $G[D_{\mathbf{8}}(Y, I, I_3) \rightarrow B' B] = -G[D_{\mathbf{8}}(Y, I, I_3) \rightarrow B B']$.

1. *Comments on the extension to $SU(4)$.*

Exact $SU(4)$ symmetry is often used to make estimates of charmed baryon couplings. This gives rise to wrong results as $SU(4)$ is badly broken. In this section we comment on the extension to $SU(4)$ of the analysis of Sec. III A [20].

The $SU(4)$ group representation includes a new quantum number charm, which comes as a new quark. The baryons fit into the product:

$$4 \otimes 4 \otimes 4 = 4^* \oplus 20' \oplus 20' \oplus 20, \quad (26)$$

the $J^P = (1/2)^+$ baryons are assigned to the $20'$ representations and the $J^P = (3/2)^+$ resonances are placed in the 20 .

In Sec. III A the decays of the dibaryon octet D_8 into an antisymmetric final state $B_8 + B_8$ were studied. Correspondingly, for charmed dibaryons the decays of the 64-plet, D_{64} , into an antisymmetric final state $B_{20'} + B_{20'}$ have to be considered, as can be seen from the decomposition:

$$20' \otimes 20' = 6 \oplus 10 \oplus 10^* \oplus 50 \oplus 64_A \oplus 64_S \oplus 70 \oplus 126. \quad (27)$$

For exact $SU(4)$ the Yukawa interaction is characterized by only one coupling constant, since the antisymmetrical 64_A representation appears only once in the decomposition of $20' \otimes 20'$.

For $SU(4)$ breaking it can be assumed, by using the spurion method, that the symmetry breaking operator transforms like the $Y = I = 0$ component of the $SU(4)$ 15 representation. Denote the symmetry breaking interaction as a spurion, S_{15} , the $(Y, I) = (0, 0)$ state of the $SU(4)$ 15 , and consider the reaction

$$S_{15} + D_{64} \rightarrow B_{20'} + B_{20'} \quad (28)$$

as $SU(4)$ invariant. Consider the decomposition

$$15 \otimes 64 = 6 \oplus 10 \oplus 10^* \oplus 50 \oplus 3(64) \oplus 70 \oplus 70^* \oplus 126 \oplus 126^* \oplus 300, \quad (29)$$

and the decomposition of $20' \otimes 20'$, Eq. (27), where the representations 6 , 50 , 64_A , and 70 are antisymmetric. For the two final baryon $20'$ be in a flavor antisymmetric state, we find that there are six $SU(4)$ breaking terms to first order, namely, $\langle 6|6 \rangle$, $\langle 50|50 \rangle$, $\langle 64|64_A \rangle$,

$\langle 64' | 64_A \rangle$, $\langle 64'' | 64_A \rangle$, and $\langle 70 | 70 \rangle$. Sum rules for SU(4) breaking along with exact SU(3) could be calculated at this first stage. We plan to present these elsewhere.

B. Decays of octet dibaryon into $B_8 + B_8$ symmetric final state.

Now we consider decays of the dibaryon octet D_8 into two baryon octets flavor symmetric final state. In principle, with λ_8 first order breaking of SU(3) there are five additional parameters g'_i ($i = 1, 2, 3, 4, 5$) in addition to g_8^0 :

$$g_8^0 \text{Tr} \{ \overline{D}_8 [B_8 B'_8 + B'_8 B_8] \}, \quad (30)$$

$$g'_1 \text{Tr} \{ \overline{D}_8 \lambda_8 [B_8 B'_8 + B'_8 B_8] \}, \quad (31)$$

$$g'_2 \text{Tr} \{ \overline{D}_8 [B_8 \lambda_8 B'_8 + B'_8 \lambda_8 B_8] \}, \quad (32)$$

$$g'_3 \text{Tr} \{ \overline{D}_8 [B_8 B'_8 + B'_8 B_8] \lambda_8 \}, \quad (33)$$

$$g'_4 \{ \text{Tr} [\overline{D}_8 B_8] \text{Tr} [B'_8 \lambda_8] + \text{Tr} [\overline{D}_{8A} B'_8] \text{Tr} [B_8 \lambda_8] \}. \quad (34)$$

$$g'_5 \text{Tr} [\overline{D}_8 \lambda_8] \text{Tr} [B_8 B'_8]. \quad (35)$$

However, the terms in Eqs. (30)-(35) are not linearly independent, there is the mathematical identity:

$$\begin{aligned} & \text{Tr} \{ \overline{D}_8 \lambda_8 [B_8 B'_8 + B'_8 B_8] \} + \text{Tr} \{ \overline{D}_8 [B_8 \lambda_8 B'_8 + B'_8 \lambda_8 B_8] \} \\ & + \text{Tr} \{ \overline{D}_8 [B_8 B'_8 + B'_8 B_8] \lambda_8 \} = \\ & \text{Tr} [\overline{D}_8 B_8] \text{Tr} [B'_8 \lambda_8] + \text{Tr} [\overline{D}_8 B'_8] \text{Tr} [B_8 \lambda_8] + \text{Tr} [\overline{D}_8 \lambda_8] \text{Tr} [B_8 B'_8]. \end{aligned} \quad (36)$$

Using (36) to eliminate the g'_2 term we finally obtain:

$$g_8^0 \text{Tr} \{ \overline{D}_8 [B_8 B'_8 + B'_8 B_8] \}, \quad (37)$$

$$g_1 \text{Tr} \{ \overline{D}_8 \lambda_8 [B_8 B'_8 + B'_8 B_8] \}, \quad (38)$$

$$g_2 \text{Tr} \{ \overline{D}_8 [B_8 B'_8 + B'_8 B_8] \lambda_8 \}, \quad (39)$$

$$g_3 \{ \text{Tr} [\overline{D}_8 B_8] \text{Tr} [B'_8 \lambda_8] + \text{Tr} [\overline{D}_{8s} B'_8] \text{Tr} [B_8 \lambda_8] \}. \quad (40)$$

$$g_4 \text{Tr} [\overline{D}_8 \lambda_8] \text{Tr} [B_8 B'_8]. \quad (41)$$

As described at the beginning of this section, we have five parameters in this case: g_8^0 from exact SU(3) and g_i ($i = 1, 2, 3, 4$) from λ_8 first order SU(3) symmetry breaking.

There are nine independent coupling constants to be determined, we take these to be:

$$G[D_8(1, 1/2, +1/2) \rightarrow p \Sigma^{0'}], \quad G[D_8(1, 1/2, +1/2) \rightarrow p \Lambda'], \quad (42)$$

$$G[D_8(0, 1, +1) \rightarrow p \Xi^{0'}], \quad G[D_8(0, 1, +1) \rightarrow \Sigma^+ \Lambda'], \quad (43)$$

$$G[D_8(0, 0, 0) \rightarrow p \Xi^{-'}], \quad G[D_8(0, 0, 0) \rightarrow \Sigma^+ \Sigma^{-'}], \quad G[D_8(0, 0, 0) \rightarrow \Lambda \Lambda'], \quad (44)$$

$$G[D_8(-1, 1/2, +1/2) \rightarrow \Sigma^+ \Xi^{-'}], \quad G[D_8(-1, 1/2, +1/2) \rightarrow \Lambda \Xi^{0'}]. \quad (45)$$

The remaining decay coupling constants are given by the same relations, Eqs. (15)-(22), with $G[D_8(0, 1, +1) \rightarrow \Sigma^+ \Sigma^{0'}] = 0$ and $G[D_8(0, 0, 0) \rightarrow \Sigma^+ \Sigma^{-'}] = -G[D_8(0, 0, 0) \rightarrow \Sigma^0 \Sigma^{0'}]$.

The nine coupling constants (42)-(45) are described by the five parameters g_8^0 and g_i ($i = 1, 2, 3, 4$). Thus, four SU(3) broken sum rules between the decay couplings constants may be deduced:

$$\begin{aligned} \sqrt{6} G[D_8(1, 1/2, +1/2) \rightarrow p \Lambda'] &= -\sqrt{2} G[D_8(1, 1/2, +1/2) \rightarrow p \Sigma^{0'}] \\ &- \sqrt{6} G[D_8(-1, 1/2, +1/2) \rightarrow \Lambda \Xi^{0'}] + G[D_8(-1, 1/2, +1/2) \rightarrow \Sigma^+ \Xi^{-'}], \end{aligned} \quad (46)$$

$$\begin{aligned} \sqrt{6} G[D_{\mathbf{8}}(0, 1, +1) \rightarrow \Sigma^+ \Lambda'] &= -2 G[D_{\mathbf{8}}(0, 1, +1) \rightarrow p \Xi^{0'}] \\ &- \sqrt{6} G[D_{\mathbf{8}}(-1, 1/2, +1/2) \rightarrow \Lambda \Xi^{0'}] + G[D_{\mathbf{8}}(-1, 1/2, +1/2) \rightarrow \Sigma^+ \Xi^{-'}], \end{aligned} \quad (47)$$

$$\begin{aligned} \sqrt{6} G[D_{\mathbf{8}}(0, 0, 0) \rightarrow p \Xi^{-'}] &= 2\sqrt{2} G[D_{\mathbf{8}}(1, 1/2, +1/2) \rightarrow p \Sigma^{0'}] \\ &+ G[D_{\mathbf{8}}(0, 1, +1) \rightarrow p \Xi^{0'}] + \sqrt{6} G[D_{\mathbf{8}}(0, 0, 0) \rightarrow \Sigma^+ \Sigma^{-'}] \\ &+ 2 G[D_{\mathbf{8}}(-1, 1/2, +1/2) \rightarrow \Sigma^+ \Xi^{-'}], \end{aligned} \quad (48)$$

$$\begin{aligned} 3\sqrt{3} G[D_{\mathbf{8}}(0, 0, 0) \rightarrow \Lambda \Lambda'] &= -8 G[D_{\mathbf{8}}(1, 1/2, +1/2) \rightarrow p \Sigma^{0'}] \\ &- 2\sqrt{2} G[D_{\mathbf{8}}(0, 1, +1) \rightarrow p \Xi^{0'}] - 3\sqrt{3} G[D_{\mathbf{8}}(0, 0, 0) \rightarrow \Sigma^+ \Sigma^{-'}] \\ &- 6\sqrt{3} G[D_{\mathbf{8}}(-1, 1/2, +1/2) \rightarrow \Lambda \Xi^{0'}] - \sqrt{2} G[D_{\mathbf{8}}(-1, 1/2, +1/2) \rightarrow \Sigma^+ \Xi^{-'}]. \end{aligned} \quad (49)$$

With identical relationships for $G[D_{\mathbf{8}}(Y, I, I_3) \rightarrow B' B] = G[D_{\mathbf{8}}(Y, I, I_3) \rightarrow B B']$.

IV. CONCLUDING REMARKS

Earlier, decays of sextaquark dibaryon decuplets ($\mathbf{10}^*$ and $\mathbf{10}$ representations) into a flavor antisymmetric final state of two ordinary baryon octets were considered [12]. In this paper we have given sum rules with λ_8 first order SU(3) symmetry breaking for decays of dibaryon octet $D_{\mathbf{8}}$ into two ordinary baryon octets $B_{\mathbf{8}} + B_{\mathbf{8}}$. The sum rules given here and in earlier papers [12, 19] can be applied into $B(\mathbf{8}) + \bar{B}(\mathbf{8})$ with appropriate changes.

Very recently LHCb it has been reported that a multiquark state $Z(4430)^-$ (tetraquark) made of charm, anti-charm, down and anti-up quarks [21] exists. This opens up a whole new area of multiquark states with charm and thus an SU(4) extension of our results is called for.

Apart from the heavy ion experiments, which we mentioned in the introduction, in our view e^+e^- annihilation reaction is best suited to find multibaryon plus antimultibaryon pairs, since the signal for resonance will be a clear signal. However LHCb will also be a good place for observing these states.

Acknowledgments

V. Gupta, E. N. Polanco-Euán, and G. Sánchez-Colón would like to thank CONACyT (México) for partial support. B. A. Bambah would like to acknowledge Ms. A. Aiswarya for checking some calculations.

-
- [1] F. J. Dyson and N.-H. Xuong, *Phys. Rev. Lett.* **13**, 815 (1964); Erratum *Phys. Rev. Lett.* **14**, 339 (1965).
 - [2] R. L. Jaffe, *Phys. Rev. Lett.* **38**, 195 (1977); Erratum *Phys. Rev. Lett.* **38**, 617 (1977).
 - [3] J. Vijande, A. Valcarce and J. M. Richard, *Phys. Rev. D* **85**, 014019 (2012).
 - [4] Y. Nambu and B. Bambah, *Phys. Rev. D* **26**, 2871 (1982).
 - [5] M. Crawford and B. Bambah, *Phys. Rev. D* **29**, 1334 (1984).
 - [6] T. Inoue et al. (HAL QCD), *Phys. Rev. Lett.* **106**, 162002 (2011).
 - [7] S. R. Beane *et al.* [NPLQCD Collaboration], *Phys. Rev. D* **85**, 054511 (2012).
 - [8] S. Cho et al. [ExHIC Collaboration], *Phys. Rev. Lett.* **106**, 212001 (2011).
 - [9] P. Burikham and E. Hirunsirisawat, *Adv. High Energy Phys.* **2011**, 123184 (2011).
 - [10] M. Bashkanov, Stanley J. Brodsky and H. Clement, *Phys. Lett. B* **727**, 438 (2013).
 - [11] P. Alderson *et al.*, *Phys. Rev. Lett.* **112**, 202301 (2014).
 - [12] V. Gupta and G. Sánchez-Colón, *Mod. Phys. Lett. A* **30**, 1550010 (2015).
 - [13] B. H. Kim *et al.* [Belle Collaboration], *Phys. Rev. D* **86**, 010001 (2012).
 - [14] M. Oka, *Phys. Rev. D* **38**, 298 (1988).
 - [15] P. Carruthers. *Introduction to Unitary Symmetry*. New York: Interscience (1966).
 - [16] J. J. de Swart, *Rev. Mod. Phys.* **35**, 916 (1963).
 - [17] M. Gell-Mann, “*The Eightfold Way: A Theory of Strong Interaction Symmetry*”, California Institute of Technology Synchrotron Laboratory Report CTSL-20 (1961), unpublished.
 - [18] S. Okubo, *Prog. Theor. Phys.* **27**, 949 (1962).
 - [19] V. Gupta and V. Singh, *Phys. Rev.* **135**, B1442 (1964).
 - [20] V. Gupta, *Pramana* **6**, 259 (1976).
 - [21] R. Aaij *et al.* [LHCb Collaboration], *Phys. Rev. Lett.* **112**, no. 22, 222002 (2014).

TABLE I: The (Y, I) states $D_{\mathbf{8}}(Y, I)$ in the $D_{\mathbf{8}}$ dibaryon octet that one obtains in the reduction $\mathbf{8} \otimes \mathbf{8}$ of two ordinary baryon octets $B_{\mathbf{8}} + B_{\mathbf{8}}$. For the SU(3) Clebsch-Gordan coefficients see Refs. [15, 16]. The average mass (in MeV) is given in the last column.

$D_{\mathbf{8}}(Y, I)$	$B_{\mathbf{8}} + B_{\mathbf{8}}$	Average mass
$D_{\mathbf{8}}(1, \frac{1}{2})$	$\frac{1}{2}[(\Sigma N')_{\frac{1}{2}} + (N \Sigma')_{\frac{1}{2}} + (N \Lambda')_{\frac{1}{2}} - (\Lambda N')_{\frac{1}{2}}]$	2093.5
$D_{\mathbf{8}}(0, 1)$	$\frac{1}{\sqrt{6}}[2(\Sigma \Sigma')_1 + (N \Xi')_1 - (\Xi N')_1]$	2514.9
$D_{\mathbf{8}}(0, 0)$	$\frac{1}{\sqrt{2}}[(N \Xi')_0 + (\Xi N')_0]$	2251.5
$D_{\mathbf{8}}(-1, \frac{1}{2})$	$\frac{1}{\sqrt{2}}[(\Sigma \Xi')_{\frac{1}{2}} + (\Xi \Sigma')_{\frac{1}{2}} + (\Lambda \Xi')_{\frac{1}{2}} - (\Xi \Lambda')_{\frac{1}{2}}]$	2472.5